

## Posterior Stabilization/Bankart Repair

### Phase I (1 – 5 days post-op)

- Wound care: monitor surgical site
- Modalities: prn for pain and inflammation (ice, IFC)
- Sling: to be worn continuously, except in therapy or during exercise sessions
- ROM: Elbow, wrist, and hand
- Exercises: None

### Phase II (5 days – 4 weeks post-op)

- Wound care: monitor site/scar management techniques
- Modalities: prn for pain and inflammation (ice, IFC)
- Sling: to be worn continuously, except in therapy or during exercise sessions
- PROM:
  - Flexion to 90 degrees
  - Abduction to 90 degrees
  - ER at 45 degrees of abduction to 30 degrees
  - IR at 45 degrees of abduction to 15-25 degrees
- Strengthening:
  - Elbow, wrist, and hand AROM
  - Fitness exercises limited to recumbent bike
  - GH joint mobilizations grade I/II for pain control

### Phase III (4 weeks – 10 weeks post-op)

- Sling:
  - At 4 weeks, D/C use at home
  - Until 6 weeks, must continue to wear sling outdoors or in public settings. D/C sling at 6 weeks
- ROM:
  - At 4-6 weeks:
    - Progress AAROM
    - IR at 45 degrees abduction to 35 degrees
  - At 6-10 weeks:
    - By 10 weeks, progress AROM/PROM to WNLs for patient, except IR at 45 degrees abduction to 35 degrees

- Strengthening:
  - At 4-6 weeks:
    - Isometric strengthening with elbow at 90 degrees and arm at side
    - Scapular stabilization exercises
  - At 6-10 weeks:
    - Light theraband at 0 degrees of abduction
    - Progress to resisted strengthening with dumbbells
    - Initiate body blade and rhythmic stabilization

#### **Phase IV (10+ weeks post-op)**

- ROM:
  - At 12 weeks,
    - Start full IR ROM and capsular stretching
    - Goal of full AROM in all planes
- Strengthening:
  - Advance as tolerated all shoulder musculature
  - Can include plyometric and proprioceptive training routines
    - At 10 weeks, 2 handed plyometrics
    - At 12 weeks, progress to single handed plyometrics

#### **Phase V (16+ weeks post-op)**

- Athletes can begin a return to throwing program (contact physician before beginning)
- Gradual resumption of supervised specific exercise
- Return to non-contact sports possible for some athletes by 3 months
- Contact/collision sport after 6 months if athlete is compliant
- Max medical improvement for athletic activities by 12 months post-op
- No weight training until 8 months post-op

Adapted From:

- 1) Brotzman SB, Wilk KE. Clinical Orthopedic Rehabilitation Second Edition. Philadelphia: Mosby 2003.