

## **Prescription for Supplies for Dressing Change**

Kelly Holtkamp, MD

Sterile Saline (eye contacts solution) Distilled water Betadine (Povidone-Iodine) Hydrogen Peroxide 4x4 Gauze Pads 2x2 Gauze Pads	<ul> <li>2-inch Kling</li> <li>3-inch Kling</li> <li>Kerlix</li> <li>Coban: 1in, 2in</li> <li>Disposable Sterile Scissors</li> <li>Gauze Non-Stick Pads</li> </ul>	<ul> <li>Adaptic (Non-Stick Dressing)</li> <li>Ace Bandage: 2in, 3in, 4in</li> <li>Disposable Gloves</li> <li>Xeroform</li> </ul>
Quantity for weeks.		
Refill PRN May Substitute		
Kelly A. Holtkamp, M.D.		Date
ressing Instructions: ret-To-Dry Dressing Change Wet the gauze with sterile saline until it is moist Place the gauze over the exposed tissue Cover the moist gauze with dry gauze Secure the dressing with the Kling or Kerlix Change the dressing once a day The moist gauze will dry and adhere to the tissue. When the gauze is removed, it will pull off the dead tissue. Sta the outer edge of the wound. You may see some bleeding, this is normal. If it is too painful, then soak the gauze sterile saline to make it easier to remove.		
<ul> <li>Dry Dressing Change</li> <li>1. Cover the wound with gauze OR non-adherent material (Xeroform, Adaptic)</li> <li>2. Secure the gauze with Kling or Kerlix</li> <li>3. Change the dressing once a day and as needed</li> </ul>		
<ol> <li>Betadine Soaking</li> <li>Pour sterile saline into a clean cont of tea.</li> <li>Place the exposed wound into the solution</li> <li>Cover the wound with gauze</li> <li>Secure the gauze with Kling or Ker</li> </ol>	fluid for 5 minutes.	of the Betadine solution to make the fluid the color

5. Perform the soak once a day.

## Signs of Infection: Notify your doctor if you experience fever, chills, night sweats, red streaking up your arm, increased swelling, redness or pain.