

Total Shoulder Arthroplasty

Phase I (1 – 5 days post-op)

- Modalities: PRN for pain and inflammation (ice, IFC)
- Sling: Ultrasling worn continuously except in therapy or during exercise sessions
- ROM:
 - Shoulder pendulums
 - o AROM: forearm, wrist, and hand
- Exercises:
 - Parascapular strengthening and scapular stabilization
 - Grip strengthening

Phase II (5 days – 4 weeks post-op)

- Wound care: Monitor site/scar management techniques
- Modalities: PRN for pain and inflammation (ice, IFC)
- Sling: Until 4 weeks post-op: Ultrasling worn continuously except in therapy or during exercise sessions
- ROM:
 - PROM progressing to AAROM progressing to AROM gradually
 - Pulleys into flexion only
 - Maximums:
 - Abduction 75 degrees
 - Flexion 120 degrees
 - ER 40 degrees (NO PROM)
 - NO ACTIVE IR
 - NO ACTIVE EXTENSION
 - AROM forearm, wrist, and hand
- Strengthening:
 - Continue periscapular strengthening and grip strengthening
 - Starting at 10 days post-op, begin isometrics for abd/ER
 - Until 8 weeks, Avoid resisted elbow flexion and supination (bicep)

Initiation Date: 06-07-04 Revised Date: 09-24-14, 3-26-19

Phone (815) 484-6990 * Fax (815) 484-6961



Phase III (4 weeks - 10 weeks post-op)

- Modalities: PRN for pain and inflammation (ice, IFC)
- Sling:
 - At 4 weeks, begin weaning out of Ultrasling in controlled environments and remove abd pillow
 - Until 6 weeks, continue to wear sling outdoors or in public settings
- ROM:
 - PROM & stretching:
 - Beginning at 4 weeks post-op, PROM stretching into flexion and scaption
 - Pulleys and table stretches
 - NO PASSIVE ER
 - Until 6 weeks, NO ACTIVE IR
 - Until 6 weeks, NO ACTIVE EXTENSION
 - AAROM/AROM:
 - By 6 weeks, progress ER to 60 degrees
 - At 6 weeks, begin ACTIVE IR and Extension
 - By 10 weeks, achieve 140-150 degrees of active flexion and scaption
- Strengthening:
 - At 6 weeks, begin Isometrics for flexion
 - At 8 weeks, begin PREs for flexion, scaption, and ER

Phase IV (10+ weeks post-op)

- ROM: Unrestricted
- Mobilizations: GH joint mobilizations
- Strengthening:
 - Progress RTC/periscapular strengthening to return to ADLs or work activity

Adapted From:

- 1) Wilk KE, Reinold MM, Andrews, JR. Total Shoulder Replacement Post-Operative Rehabilitation Program. Winchester MA; Advanced Continuing Education Institute, 2019.
- 2) The Brigham and Women's Hospital Inc. Dept. of Rehabilitation Services, 2016.

Initiation Date: 06-07-04 Revised Date: 09-24-14, 3-26-19

Phone (815) 484-6990 * Fax (815) 484-6961